



Congenital Cardiology Solutions

REAL TIME THREE DIMENSIONAL TRANSESOPHAGEAL ECHOCARDIOGRAPHY COUPLED WITH DIAGNOSTIC CATHETERIZATION TO DETERMINE LEFT VENTRICULAR PRESSURE-VOLUME RELATIONSHIPS AND MYOCONTRACTILITY

ACC Moderated Poster Contributions
McCormick Place South, Hall A
Sunday, March 25, 2012, 11:00 a.m.-Noon

Session Title: Congenital Cardiology Solutions: Roles of Imaging
Abstract Category: 27. Congenital Cardiology Solutions: Pediatric
Presentation Number: 1134-49

Authors: Shelby Kutty, Sarah Nordmeyer, Asif Padiyath, Paul Gribben, Ling Li, Shunji Gao, Sheela Rangamani, Lucas Drvol, John Lof, David Danford, Titus Kuehne, University of Nebraska Medical Center and Children's Hospital and Medical Center, Omaha, NE, USA, Department of Congenital Heart Disease/Pediatric Cardiology, Deutsches Herzzentrum Berlin, Berlin, Germany

Background: We investigated left ventricular (LV) pressure volume relationships (PVR) and myocontractility indexes using real time three-dimensional transesophageal echocardiography (RT3DTEE) and conductance catheterization (CC).

Methods: CC used 6 F catheters in the LV via right carotid cut-down (Millar, MPVS Ultra) in pigs. Simultaneous CC and RT3DTEE (single beat, X7-2 X matrix, iE33, Philips) data were obtained with the animal ventilated (end-expiration), and paralyzed in 3 conditions (3 measurements in each): baseline, inferior vena caval (IVC) occlusion, and IVC occlusion during dobutamine. CC derived LV pressures (Labchart) were integrated with RT3DTEE volumes (QLab) on a beat-to-beat basis using ECG trigger signal. From CC and RT3DTEE PV loops, end systolic PVR (Emax) were determined.

Results: In 6 pigs studied, there were excellent agreements between the RT3DTEE and CC for parameters derived at baseline and with dobutamine. At baseline, the mean \pm SD (mmHg/mL) were Emax-CC 1.86 ± 1.1 and Emax-RT3DTEE 1.78 ± 1.2 ($p=0.50$). On dobutamine, mean Emax-CC was 3.43 ± 1.5 and Emax-RT3DTEE 3.60 ± 1.23 ($p=0.17$). Emax normalized to end diastolic volumes were 0.034 ± 0.02 for CC and 0.033 ± 0.02 for RT3DTEE at baseline ($p=0.43$); these were 0.081 ± 0.04 and 0.084 ± 0.03 respectively on dobutamine ($p=0.13$).

Conclusion: Emax can be determined from RT3DTEE, and agrees well with CC. Derivation of PVR and Emax using RT3DTEE permits integration of intrinsic myocardial functional parameters in the clinical context.

